

EVOLUTION OF LITHIUM IN THE MILKY WAY

D. ROMANO

*International School for Advanced Studies, SISSA/ISAS, Via Beirut 2-4,
34014 Trieste, Italy
E-mail: romano@sissa.it*

F. MATTEUCCI

*Dipartimento di Astronomia, Università di Trieste, Via G.B. Tiepolo 11,
34131 Trieste, Italy
E-mail: matteucci@ts.astro.it*

P. VENTURA, F. D'ANTONA

*Osservatorio Astronomico di Roma, Via Frascati 33,
00040 Monte Porzio Catone, Italy
E-mail: paolo, dantona@coma.mporzio.astro.it*

We adopt up-to-date ${}^7\text{Li}$ yields from asymptotic giant branch (AGB) stars in order to study the temporal evolution of this element in the solar neighbourhood. Several lithium stellar sources are considered besides the AGBs: Type II supernovae (SNe), novae, low-mass giants. Galactic cosmic ray (GCR) nucleosynthesis is taken into account as well. We conclude that AGB stars do not substantially contribute to ${}^7\text{Li}$ enrichment on a Galactic scale. Therefore, a significant ${}^7\text{Li}$ production from novae and low-mass stars is needed to explain the late, steep rise of the ${}^7\text{Li}$ abundance in disk stars and the meteoritic ${}^7\text{Li}$ abundance.

1 Model Prescriptions and Results

The upper envelope of the observed $\log \epsilon({}^7\text{Li})$ vs. $[\text{Fe}/\text{H}]$ diagram can constrain models of chemical evolution, provided that it actually traces the ${}^7\text{Li}$ enrichment history of the interstellar medium (ISM) in the solar neighbourhood. A nearly flat plateau at low metallicities (the *Spite Plateau*) is followed by a steep rise afterwards (big dots in Figure 1): these features are interpreted as due to ${}^7\text{Li}$ enhancement from a lower primordial value to a higher present one. The main purpose of this contribution is to show how the interplay between several categories of ${}^7\text{Li}$ producers may explain the rise from the Spite plateau and the meteoritic ${}^7\text{Li}$ abundance.

The adopted model assumes that the Galaxy forms out of two main episodes of accretion: the first one builds up the halo, the second one the disk. The Sun is located at a distance of 8 kpc from the Galactic center, the adopted Galactic age is 14 Gyr. Details on the model parameters and basic equations can be found in Chiappini *et al.*^{1,2}; detailed ${}^7\text{Li}$ synthesis prescriptions are given in Romano *et al.*³.




Figure 1: Model predictions compared to the data. The contribution from each category of stellar ${}^7\text{Li}$ producers is shown in the left panel (see text for details); the ${}^7\text{Li}$ evolution obtained by adding together in the model all the stellar sources and GCR nucleosynthesis is shown in the right panel for three different values of the primordial abundance. The results at $[\text{Fe}/\text{H}] > -0.5$ dex are nearly the same, owing to the consumption of ${}^7\text{Li}$ by stellar astration during the Galactic lifetime.

In Figure 1, left panel, the contributions to ${}^7\text{Li}$ enrichment from different stellar sources are shown: *i*) AGB stars (*short-dashed line*); *ii*) Type II SNe (*dotted line*); *iii*) low-mass giants (*long-dashed line*); and *iv*) novae (*continuous line*). Model predictions in the right panel of Figure 1 are obtained by taking into account in the model all these stellar sources plus *v*) GCRs.

The main conclusions are that AGBs are not substantial ${}^7\text{Li}$ producers on a Galactic scale (see also the contributions by P. Ventura and F. D’Antona, these proceedings) and that the meteoritic ${}^7\text{Li}$ abundance can be reproduced only if other stellar sources acting at late times (novae and low-mass stars) are contributing a substantial fraction ($\sim 60\%$ in our model) of the meteoritic ${}^7\text{Li}$. However, it should be said that the fractional contribution of each of these two sources is still very uncertain, mainly due to uncertainties in the evolutionary histories of the progenitor stars and in the stellar yields.

References

1. C. Chiappini *et al*, *ApJ* **477**, 765 (1997).
2. C. Chiappini *et al*, *ApJ*, accepted (2001).
3. D. Romano *et al*, *A&A*, submitted (2001).